

Erratum for ‘The Chalk Group (Upper Cretaceous) of the Northern Province, eastern England – a review’, *Proceedings of the Yorkshire Geological Society*, 62, 153–177



Simon F. Mitchell

Department of Geography and Geology, The University of the West Indies, Mona, Kingston 7, Jamaica
simon.mitchell@uwimona.edu.jm

Errors were found in Figures 2, 17 and 19. The correct figures are as follows.

| Stage | Traditional Northern Zones | Zones used herein | Northern P. | Southern Prov. | | |
|----------------------|----------------------------|---------------------------------|-----------------------|---------------------|--------------------------|----|
| lower Campanian | Sphenocerasmus lingua | Scaphites binodosus | Flamborough Formation | Newhaven Chalk | | |
| | | Goniotheuthis granulataquadrata | | | | |
| | | Unitacrinus anglicus | | | | |
| Santonian | U | Marsupites testudinarius | Bumham Formation | Seaford Chalk | | |
| | | Marsupites laevigatus | | | | |
| | M | Unitacrinus socialis | | | | |
| | L | Hagenowia rostrata | | | | |
| Coniacian | Hagenowia rostrata | Cordiceramus cordiformis | Bumham Formation | Lewes Nodular Chalk | | |
| | | Cladocerasmus undulatoaplicatus | | | | |
| Turonian | Micraster cortestudinarium | Micraster cortestudinarium | Welton Formation | Holywell Nodular | | |
| | Sternotaxis plana | Plesiocorys plana | | | | |
| | Terebratulina lata | Terebratulina lata | | | | |
| Cenomanian | Mytiloides spp. | Mytiloides spp. | Ferriby Formation | Zig Zag Chalk | | |
| | | Sciponeras gracile | | | Praeactinocamax plenus | |
| | | Holaster tercensis | | | Crassiholaster trecensis | |
| Albian | Belemnites minimus | Crassiholaster subglobosus | Hunstanton Formation | West Melbury | | |
| | | Neohibolites praeultimus | | | Unconformity | |
| | | Neohibolites ernsti | | | | UG |
| | | Neohibolites oxycaudatus | | | | |
| Neohibolites minimus | Gault Clay | | | | | |

Fig. 2. Lithostratigraphic nomenclature of the Chalk of the Northern Province compared with that of the Southern Province. The traditional zones for the Northern Province are shown together with those used here. UG, Upper Greensand.




Fig. 17. Distribution of selected calcitic macrofossils in the Hunstanton Formation at Hunstanton.




Fig. 19. Distribution of selected fossils in the Hunstanton Formation and lowermost Ferriby Formation at South Ferriby. Belemnite biohorizons BB2 and BB3 with *Neohibolites ultimus* in the lower Cenomanian are shown.